

Analysis of Methane Emissions in Alaska Permafrost Areas Using Sentinel-5P Data

Sheik Arief Abdaly^{1,2}

¹Department of Physics, Presidency University, Kolkata 700073, India.

²Institute of Natural Sciences and Applied Technology, Kolkata 700032, India

E-mail:sheikariefabdaly@gmail.com

Received: July 4, 2024, Revised: August 8, 2024, Accepted: August 17, 2024, Published: September 30, 2024

Abstract

Methane (CH₄) is a potent greenhouse gas with significant implications for climate change, especially in Arctic permafrost regions where thawing may release large quantities of stored methane. This study investigates the spatiotemporal variations of methane concentrations over Alaska's permafrost regions from 2019 to 2023 using Sentinel-5P satellite data. Monthly averages were derived, and statistical analyses, including seasonal decomposition and autocorrelation, were performed to reveal trends, seasonality, and the underlying dynamics. Our findings suggest an increasing methane concentration trend with distinct seasonal variations, highlighting the role of permafrost regions in the global methane cycle.

Keywords: Autocorrelation, Anthropogenic, Time series analysis

I. Introduction

Methane is the second most important anthropogenic greenhouse gas after carbon dioxide, with a global warming potential significantly higher over short timescales. Methane contributes roughly 16% to the radiative forcing from all well-mixed greenhouse gases. Current estimates of the global methane source [Kirschke, 2013] strength converge at 550–900 Tg yr⁻¹, with a contribution of 25 ± 14 Tg yr⁻¹ from high-latitude wetlands in the Arctic tundra. These wetlands and other inland waters form a globally relevant methane source and are the most important source of uncertainty in the global methane budget estimate. Reasons for this uncertainty include the shortage of flux observation sites and short time series, the high spatial heterogeneity of permafrost-affected landscapes [Schuur, 2015], the pronounced temporal dynamics of methane emissions and the complex controls on methane production, transport and oxidation. With a warming in the Arctic region that is at least twice the global average, methane emissions are projected to increase in response to permafrost degradation. Permafrost regions are already releasing additional methane due to thaw. Besides sparse indirect indications of increased carbon loss neither atmospheric measurements nor inversion or biospheric models currently provide clear trends in methane emissions from high-latitude wetlands for the recent past. It also remains unclear whether rising temperatures equally affect the two counteracting processes of methane production and oxidation. Arctic permafrost regions, such as Alaska, are vital for understanding methane emissions caused by thawing organic matter in frozen soils. These emissions significantly contribute to climate change, making their monitoring essential for predicting climate feedback loops. This study utilizes Sentinel-5P satellite data to analyze methane emissions in Alaska's permafrost regions from 2019 to 2023. Key objectives include:

1. Quantifying temporal variations in methane concentrations during this period.
2. Exploring seasonal trends and correlations to assess natural and climate-driven influences.
3. Identifying potential drivers, such as temperature anomalies, soil moisture, and human activities, accelerating methane release.

II. Materials and methods

a. Study Area

The study centers on Alaska's permafrost regions, defined by the geographical bounds of $[-163.0^{\circ}, 66.0^{\circ}][-163.0^{\circ}, 66.0^{\circ}]$ to $[-141.0^{\circ}, 69.0^{\circ}][-141.0^{\circ}, 69.0^{\circ}]$. This region encompasses extensive permafrost landscapes that are particularly vulnerable to rising global temperatures. Permafrost in these areas contains significant stores of organic carbon, which can decompose upon thawing, releasing methane (CH_4) into the atmosphere. The geographical boundaries selected for this study ensure the inclusion of diverse permafrost ecosystems, ranging from continuous to discontinuous zones. These areas are of critical importance for understanding the feedback mechanisms between permafrost thaw and climate change. The analysis leverages Sentinel-5P satellite data to monitor methane concentrations, providing detailed spatiotemporal insights. The chosen bounds also align with the region's significant contribution to Arctic carbon fluxes. By focusing on these coordinates, the study captures the dynamics of methane emissions influenced by permafrost degradation. This approach highlights the region's role in the global greenhouse gas budget. Such investigations are vital for predicting and mitigating the climate impacts of Arctic permafrost changes.

b. Data Source

Methane data were sourced from the Sentinel-5P TROPospheric Monitoring Instrument (TROPOMI), specifically the `CH4_column_volume_mixing_ratio_dry_air` product. Data spanning from January 1, 2019, to December 31, 2023, were used.

c. Data Processing

The following steps were implemented:

Data Filtering: Sentinel-5P data were filtered for the specified ROI and timeframe using Google Earth Engine (GEE).

Monthly Aggregation: Monthly average methane concentrations were computed for each year.

Statistical Analysis: Mean and standard deviation were calculated, and the temporal data were decomposed into trend, seasonal, and residual components using seasonal decomposition.

III. Analytical Methods

The following steps were implemented:

Seasonal Decomposition: Employed to isolate the trend, seasonality, and residuals in methane emissions.

Autocorrelation Analysis: Used to detect patterns and correlations over time lags.

Seasonal Comparisons: Methane concentrations were compared across years to evaluate interannual variations.

IV. Results

The image illustrates the seasonal decomposition of methane (CH_4) concentrations over Alaska's permafrost regions from 2019 to 2023, using an additive model. It comprises four panels:

a. Observed Methane Concentration: The top panel displays the original time series data, showing monthly mean methane concentrations. A noticeable upward trend and seasonal variations are evident.

b. Trend Component: The second panel isolates the long-term trend, highlighting a gradual increase in methane levels over the years, indicative of persistent emissions possibly linked to permafrost thaw.

c. Seasonal Component: The third panel captures the recurring seasonal pattern, with methane levels fluctuating annually, likely driven by temperature changes and thaw-freeze cycles in the permafrost.

d. Residuals: The bottom panel represents the irregular variations or unexplained deviations from the observed data, showing noise or anomalies not accounted for by the trend or seasonality.

This decomposition provides insights into methane emission dynamics, separating the contributions of trend, seasonality, and residual variability, essential for understanding the underlying processes in Arctic permafrost regions.




Figure 1: Components of time series analysis of methane emission for Alaska permafrost.

V. Discussion

a. Trends in Methane Concentrations

The decomposition analysis revealed a consistent upward trend in methane concentrations across the study period (2019–2023). The observed trend aligns with increasing temperatures and permafrost thawing in the region, potentially releasing methane from organic matter decomposition.

b. Seasonal Variations

Seasonal decomposition highlighted significant methane peaks during late summer and early fall (August–September). This is likely driven by:

- i. Increased microbial activity during thawing seasons.
- ii. Wetland emissions amplified by warmer temperatures.

c. Autocorrelation Analysis

Autocorrelation plots indicated moderate lagged correlations, suggesting that methane concentrations are influenced by persistent seasonal and environmental factors. The peaks at lags corresponding to 12 months reinforce the annual cyclicity of emissions.




Figure 2: The image presents the autocorrelation function (ACF) plot of monthly methane emissions in Alaska's permafrost regions from 2019 to 2023.

d. Key features include:

Lags on the X-axis: Represent the time intervals (in months) at which the correlation of the methane time series is analyzed with itself.

Autocorrelation on the Y-axis: Indicates the strength of correlation between the time series values at different lags.

Significance Boundaries: The shaded blue region represents the 95% confidence interval. Points falling outside this interval indicate statistically significant autocorrelation.

The ACF reveals notable positive autocorrelations at shorter lags (e.g., up to ~6 months), which gradually diminish as the lag increases, suggesting a seasonal pattern. Negative correlations are observed at longer lags, hinting at periodic oscillations in methane levels. This analysis supports the presence of regular seasonal fluctuations and potential trends in methane emissions linked to natural cycles in permafrost dynamics.

e. Interannual Variations

Seasonal plots by year showed year-to-year variability in methane emissions. While seasonal patterns remained consistent, the magnitude of emissions increased, particularly during warmer months, reflecting potential intensification of thawing and emissions over time.

f. Implications

Our findings underscore the importance of monitoring methane emissions in permafrost regions. The observed trends and seasonal dynamics emphasize the need for:

- Enhanced Monitoring: High-resolution, year-round satellite measurements to track permafrost methane dynamics.
- Climate Modeling: Integration of permafrost methane feedbacks into global climate models.
- Mitigation Efforts: Strategies to reduce anthropogenic warming and limit permafrost degradation.

VI. Conclusion

This study provides a comprehensive analysis of methane emissions over Alaska's permafrost regions using Sentinel-5P data. The identified trends and seasonal patterns highlight the critical role of permafrost areas in the global methane budget. Continued observation and analysis are crucial for understanding and mitigating the impacts of climate change.

Acknowledgements

The author would like to thank the Institute of Natural Sciences and Applied Technology, Kolkata for providing infrastructure facilities to carry out the research.

References

- [1] Copernicus Sentinel-5P (2024). *Sentinel-5P Product User Manual*.
- [2] Schuur, E. A. G., et al. (2015). Climate change and the permafrost carbon feedback. *Nature*.
- [3] Kirschke, S., et al. (2013). Three decades of global methane sources and sinks. *Nature Geoscience*.